
















Acetaminophen (Tylenol®) Dosing Chart

Dose: Give every 4-6 hours as needed, no more than 5 times in 24 hours. Do not use in children under 3 months of age -- unless advised by your clinician.

Weight-banding (lbs, kg)	Dosage	Liquid			Solid		
		Infant's Acetaminophen (160 mg/5mL)	Children's Acetaminophen (160 mg/5mL)	Children's Acetaminophen Dissolvable Packets (160 mg/powder pack)	Children's Acetaminophen Chewables (160 mg)	Acetaminophen Regular Strength Tablets (325 mg)	Acetaminophen Extra Strength Tablets (500 mg)
6 – 11 lbs (3 to 5 kg)	40	1.25 mL 	-	-	-	-	-
12 – 17 lbs (5 to 7 kg)	80	2.5 mL 	-	-	1/2 tablet 	-	-
18 – 23 lbs (8 to 10 kg)	120	3.75 mL 	-	-	-	-	-
24 – 35 lbs (11 to 15 kg)	160	5 mL 	5 mL → 	1 packet 	1 tablet 	-	-
36 – 47 lbs (16 to 21 kg)	240	7.5 mL (= 5 mL + 2.5 mL)  and 	7.5 mL ← 	-	-	1 1/2 tablets 	-
48 – 59 lbs (22 to 26 kg)	320 – 325	10 mL (= 5mL + 5mL)  and 	10 mL → 	2 packets 	2 tablets 	1 tablet 	--
60 – 71 lbs (27 to 32 kg)	325 – 400	-	12.5 mL ← 	2 packets 	2 1/2 tablets 	1 tablet 	-
72 – 95 lbs (33 to 43 kg)	480 – 500	-	15 mL → 	3 packets 	3 tablets 	1 1/2 tablets 	1 tablet 
96 or more lbs (≥ 44 kg)	500 – 650	-	20 mL 10 mL →  and 10 mL → 	-	-	4 tablets 	2 tablets 
							1 tablet 

Ibuprofen (Motrin®, Advil®) Dosing Chart

Dose: Give every 6-8 hours, as needed, no more than 4 times in 24 hours.
Do not use under age 6 months unless advised by your clinician.













Weight-banding (lbs, kg)	Dosage (mg)	Liquid		Solid	
		Infant's Ibuprofen Concentrated Drops (50 mg / 1.25 mL) -included syringe is marked for 0.625 mL, 1.25 mL, & 1.875 mL	Children's Ibuprofen (100 mg / 5 mL) *comes with dosage cup*	Children's Ibuprofen Chewable Tablets (100 mg)	Ibuprofen Tablets or Gel Caps (200 mg)
12-17 lbs (6 to 7 kg)	50 mg	1.25 mL 	2.5 mL ^a	½ tablet 	-
18-23 lbs (8 to 10 kg)	75 mg	1.875 mL 	4 mL ^a	-	-
24-35 lbs (11 to 15 kg)	100 mg	2.5 mL (=1.875 + 0.625 mL)  and 	5 mL → 	1 tablet 	-
36-47 lbs (16 to 21 kg)	150 mg	3.75 mL (=1.875 + 1.875 mL)  and 	7.5 mL ← 	1 ½ tablets 	-
48-59 lbs (22 to 26 kg)	200 mg	5 mL ^a	10 mL → 	2 tablets 	1 tablet 
60-71 lbs (27 to 32 kg)	250 mg	-	12.5 mL ← 	2 ½ tablets 	1 tablet 
72-95 lbs (33 to 43 kg)	300 mg	-	15 mL → 	3 tablets 	1 ½ tablets ^b 
96 lbs & over (≥ 44 kg)	400 mg	-	20 mL (=10 mL + 10 mL) 10 mL →  and 10 mL → 	4 tablets 	2 tablets 

Table graphics adapted from HealthyChildrens.org (from the American Academy of Pediatrics)

^aUse of this dosage will require an oral syringe not provided in original OTC package. Ask your retail pharmacy for assistance.

^bFor ½ tablet measurements: use solid tablets only – do NOT break open gel caps

Important Notes

1. Ask your health care provider or pharmacist which formulation is best for your child.
2. Give dose based on your child's weight. Do not give more medication than recommended.
3. Always use the included measuring device OR an oral syringe given by the pharmacy. Kitchen spoons are *not* accurate measuring devices.
4. If you have questions about dosing or any other concern, call your health care provider.